arXiv:1209.0873v1 [math.CA] 5 Sep 2012

POWER MEAN INEQUALITY OF GENERALIZED TRIGONOMETRIC FUNCTIONS

BARKAT ALI BHAYO AND MATTI VUORINEN

Abstract. We study the convexity/concavity properties of the generalized p-trigonometric functions in the sense of P. Lindqvist with respect to the power means.

2010 Mathematics Subject Classification: 33C99, 33B99
Keywords and phrases: Eigenfunctions $\sin_p$, power mean, generalized trigonometric functions, convexity, inequalities.

1. Introduction

The generalized trigonometric and hyperbolic functions depending on a parameter $p>1$ were studied by P. Lindqvist in 1995 [L]. For the case when $p=2$, these functions coincide with elementary functions. Later on numerous authors have extended this work in various directions see [BE, BEM1, BEM2, DM, LP].

For $t\in\mathbb{R}$ and $x,y>0$, the Power Mean M_t of order t is defined by

$$M_t=\begin{cases}\left(\dfrac{x^t+y^t}{2}\right)^{1/t}, & t\neq 0,\\ \sqrt{x\,y}, & t=0\,.\end{cases}$$

1.1. **Theorem.** *For $p>1$, $t\geq 0$ and $r,s\in(0,1)$, we have*

(1) $\operatorname{arcsin}_p(M_t(r,s))\leq M_t(\operatorname{arcsin}_p(r),\operatorname{arcsin}_p(s))\,,$

(2) $\operatorname{artanh}_p(M_t(r,s))\leq M_t(\operatorname{artanh}_p(r),\operatorname{artanh}_p(s))\,,$

(3) $\operatorname{arctan}_p(M_t(r,s))\geq M_t(\operatorname{arctan}_p(r),\operatorname{arctan}_p(s))\,,$

(4) $\operatorname{arsinh}_p(M_t(r,s))\geq M_t(\operatorname{arsinh}_p(r),\operatorname{arsinh}_p(s))\,.$

1.2. **Theorem.** *For $p>1$, $t\geq 1$ and $r,s\in(0,1)$, the following relations hold*

(1) $\sin_p(M_t(r,s))\geq M_t(\sin_p(r),\sin_p(s))\,,$

(2) $\cos_p(M_t(r,s))\leq M_t(\cos_p(r),\cos_p(s))\,,$

`File: PM_pfuncs120904.tex, printed: 2012-9-6, 0.39`

(3) $\tan_p(M_t(r,s)) \le M_t(\tan_p(r), \tan_p(s))\,,$

(4) $\tanh_p(M_t(r,s)) \ge M_t(\tanh_p(r), \tanh_p(s))\,,$

(5) $\sinh_p(M_t(r,s)) \le M_t(\sinh_p(r), \operatorname{arsinh}_p(s))\,.$

In [BV2], there are some results which are the special case of the above theorems when $t = 0$ and $t = 2$.

Generalized convexity/concavity with respect to general mean values has been studied recently in [AVV2].

Let $f : I \to (0,\infty)$ be continuous, where I is a subinterval of $(0,\infty)$. Let M and N be any two mean values. We say that f is MN-convex (concave) if

$$f(M(x,y)) \le (\ge) N(f(x), f(y)) \quad \text{for} \quad \text{all } x, y \in I\,.$$

In conclusion, we see that the above results are (M_t, M_t)-convexity or (M_t, M_t)-concavity properties of the functions involved. In view of [AVV2], it is natural to expect that similar results might also hold for some other pairs (M, N) of mean values.

2. PRELIMINARIES

We introduce some notation and terminology for the satatement of the main results.

Given complex numbers a, b and c with $c \neq 0, -1, -2, \ldots$, the *Gaussian hypergeometric function* is the analytic continuation to the slit place $\mathbb{C} \setminus [1,\infty)$ of the series

$$F(a,b;c;z) = {}_2F_1(a,b;c;z) = \sum_{n=0}^{\infty} \frac{(a,n)(b,n)}{(c,n)} \frac{z^n}{n!}, \qquad |z| < 1.$$

Here $(a,0) = 1$ for $a \neq 0$, and (a,n) is the *shifted factorial function* or the *Appell symbol*

$$(a,n) = a(a+1)(a+2)\cdots(a+n-1)$$

for $n \in \mathbb{Z}_+$, see [AS].

Lets start the discussion of eigenfunctions of one-dimensional p-Laplacian Δ_p on $(0,1)$, $p \in (1,\infty)$. The eigenvalue problem [DM]

$$-\Delta_p u = -\left(|u^{'}|^{p-2}u^{'}\right)^{'} = \lambda |u|^{p-2}u, \quad u(0) = u(1) = 0,$$

has eigenvalues

$$\lambda_n = (p-1)(n\pi_p)^p,$$

and eigenfunctions

$$\sin_p(n\pi_p\, t), \quad n \in \mathbb{N},$$

where $\sin_p$ is the inverse function of $\arcsin_p$, which is defined below, and

$$\pi_p = \frac{2}{p}\int_0^1 (1-s)^{-1/p} s^{1/p-1} ds = \frac{2}{p} B\left(1 - \frac{1}{p}, \frac{1}{p}\right) = \frac{2\pi}{p\,\sin(\pi/p)}\,,$$

with $\pi_2 = \pi$.

Lets consider the following homeomorphisms

$$\sin_p : (0, a_p) \to I, \quad \cos_p : (0, a_p) \to I, \quad \tan_p : (0, b_p) \to I,$$
$$\sinh_p : (0, c_p) \to I, \quad \tanh_p : (0, \infty) \to I\,,$$

where $I = (0, 1)$ and

$$a_p = \frac{\pi_p}{2},\ b_p = 2^{-1/p} F\left(\frac{1}{p}, \frac{1}{p}; 1 + \frac{1}{p}; \frac{1}{2}\right),\ c_p = F\left(1\,, \frac{1}{p}; 1 + \frac{1}{p}\,; 1\right).$$

For $x \in I$, their inverse functions are defined as

$$\begin{aligned}
\operatorname{arcsin}_p x &= \int_0^x (1 - t^p)^{-1/p} dt = x\, F\left(\frac{1}{p}, \frac{1}{p}; 1 + \frac{1}{p}; x^p\right)\\
&= x(1 - x^p)^{(p-1)/p} F\left(1, 1; 1 + \frac{1}{p}; x^p\right)\,,\\
\operatorname{arctan}_p x &= \int_0^x (1 + t^p)^{-1} dt = x\, F\left(1, \frac{1}{p}; 1 + \frac{1}{p}; -x^p\right)\\
&= \left(\frac{x^p}{1 + x^p}\right)^{1/p} F\left(\frac{1}{p}, \frac{1}{p}; 1 + \frac{1}{p}; \frac{x^p}{1 + x^p}\right)\,,\\
\operatorname{arsinh}_p x &= \int_0^x (1 + t^p)^{-1/p} dt = x\, F\left(\frac{1}{p}\,, \frac{1}{p}; 1 + \frac{1}{p}; -x^p\right)\\
&= \left(\frac{x^p}{1 + x^p}\right)^{1/p} F\left(1, \frac{1}{p}; 1 + \frac{1}{p}; \frac{x^p}{1 + x^p}\right)\,,\\
\operatorname{artanh}_p x &= \int_0^x (1 - t^p)^{-1} dt = x\, F\left(1\,, \frac{1}{p}; 1 + \frac{1}{p}; x^p\right)\,,
\end{aligned}$$

and by [BE, Prop 2.2] $\operatorname{arccos}_p x = \operatorname{arcsin}_p((1 - x^p)^{1/p})$. In particular, they reduce to the familiar functions for the case $p = 2$.

The above functions were generalized in two parameters (p, q) in [BV2, T, EGL].

For $x \in I = [0, 1]$

$$\operatorname{arcsin}_{p,q} x = \int_0^x (1 - t^q)^{-1/p} dt = x\, F\left(\frac{1}{p}, \frac{1}{q}; 1 + \frac{1}{p}; x^q\right).$$

We also define $\operatorname{arccos}_{p,q} x = \operatorname{arcsin}_{p,q}((1 - x^p)^{1/q})$ (see [EGL, Prop. 3.1]), and

$$\operatorname{arsinh}_{p,q} x = \int_0^x (1 + t^q)^{-1/p} dt = x\, F\left(\frac{1}{p}, \frac{1}{q}; 1 + \frac{1}{q}; -x^q\right).$$

Their inverse functions are

$$\sin_{p,q} : (0, \pi_{p,q}/2) \to (0, 1), \quad \cos_{p,q} : (0, \pi_{p,q}/2) \to (0, 1),$$
$$\sinh_{p,q} : (0, n_{p,q}) \to (0, 1), \quad n_{p,q} = \frac{1}{2^{1/p}} F\left(1, \frac{1}{p}; 1 + \frac{1}{q}; \frac{1}{2}\right).$$

For easy reference we record the following lemma from [AVV1], sometimes which is called the *monotone l'Hospital rule.*

2.1. **Lemma.** [AVV1, Theorem 1.25] *For* $-\infty < a < b < \infty$*, let* $f, g : [a, b] \to \mathbb{R}$ *be continuous on* $[a, b]$*, and be differentiable on* (a, b)*. Let* $g'(x) \neq 0$ *on* (a, b)*. If* $f'(x)/g'(x)$ *is increasing (decreasing) on* (a, b)*, then so are*

$$[f(x) - f(a)]/[g(x) - g(a)] \quad \textit{and} \quad [f(x) - f(b)]/[g(x) - g(b)].$$

If $f'(x)/g'(x)$ *is strictly monotone, then the monotonicity in the conclusion is also strict.*

For the next two lemmas see [BV1, Theorems 1.1, 1.2, 2.5 & Lemma 3.6].

2.2. **Lemma.** *For* $p > 1$ *and* $x \in (0, 1)$*, we have*

(1) $\left(1 + \dfrac{x^p}{p(1+p)}\right) x < \operatorname{arcsin}_p x < \dfrac{\pi_p}{2}\, x,$

(2) $\left(1 + \dfrac{1 - x^p}{p(1+p)}\right) (1 - x^p)^{1/p} < \operatorname{arccos}_p x < \dfrac{\pi_p}{2}\, (1 - x^p)^{1/p},$

(3) $\dfrac{(p(1+p)(1+x^p) + x^p)x}{p(1+p)(1+x^p)^{1+1/p}} < \operatorname{arctan}_p x < 2^{1/p}\, b_p \left(\dfrac{x^p}{1+x^p}\right)^{1/p}.$

(4) $z\left(1 + \dfrac{\log(1+x^p)}{1+p}\right) < \operatorname{arsinh}_p x < z\left(1 + \frac{1}{p}\log(1+x^p)\right), \ z = \left(\dfrac{x^p}{1+x^p}\right)^{1/p},$

(5) $x\left(1 - \dfrac{1}{1+p}\log(1-x^p)\right) < \operatorname{artanh}_p x < x\left(1 - \frac{1}{p}\log(1-x^p)\right).$

2.3. **Lemma.** *For* $p, q > 1$ *and* $r, s \in (0, 1)$*, the following inequalities hold:*

(1) $\operatorname{arcsin}_p(\sqrt{r\, s}) \leq \sqrt{\operatorname{arcsin}_p(r) \operatorname{arcsin}_p(s)}\,,$

(2) $\operatorname{artanh}_p(\sqrt{r\, s}) \leq \sqrt{\operatorname{artanh}_p(r) \operatorname{artanh}_p(s)}\,,$

(3) $\sqrt{\operatorname{arsinh}_p(r) \operatorname{arsinh}_p(s)} \leq \operatorname{arsinh}_p(\sqrt{r\, s})\,,$

(4) $\sqrt{\operatorname{arctan}_p(r) \operatorname{arctan}_p(s)} \leq \operatorname{arctan}_p(\sqrt{r\, s})\,,$

(5) $\pi_{\sqrt{pq}} \leq \sqrt{\pi_p\, \pi_q}.$

2.4. **Lemma.** *For* $m \geq -1$*,* $p > 1$*, the following functions*

(1) $f_1(x) = \left(\dfrac{\operatorname{arcsin}_p x}{x}\right)^m \dfrac{d}{dx}(\operatorname{arcsin}_p x),$

(2) $f_2(x) = \left(\dfrac{\operatorname{artanh}_p x}{x}\right)^m \dfrac{d}{dx}(\operatorname{artanh}_p x),$

are increasing in $x \in (0, 1)$*, and*

(3) $f_3(x) = \left(\dfrac{\operatorname{arctan}_p x}{x}\right)^m \dfrac{d}{dx}(\operatorname{arctan}_p x)$,

(4) $f_4(x) = \left(\dfrac{\operatorname{arsinh}_p x}{x}\right)^m \dfrac{d}{dx}(\operatorname{arsinh}_p x)$,

are decreasing in $x \in (0,1)$.

Proof. By definition

$$f_1(x) = \left(\frac{\arcsin_p x}{x}\right)^m \frac{1}{(1-x^p)^{1/p}}.$$

For $m \geq 0$, $\left(\frac{\arcsin_p x}{x}\right)^m$ is increasing by Lemma 2.1, and clearly $(1-x^p)^{1/p}$ is increasing. For the case $m \in [-1,0)$, we define

$$h_1(x) = \left(\frac{x}{\arcsin_p x}\right)^s \frac{1}{(1-x^p)^{1/p}}, \quad s \in (0,1].$$

We get

$$\begin{aligned} h_1'(x) &= \xi((1-x^p)^{1/p}(x^p + s(1-x^p))F_1(x) - s(1-x^p)) \\ &> \xi\left((1-x^p)^{1/p}(x^p + s(1-x^p))(1 + \frac{x^p}{p(1+p)}) - s(1-x^p)\right) > 0, \end{aligned}$$

by Lemma 2.2(1), where

$$\xi = \frac{(1-x^p)^{-(1+2/p)}}{x}\left(\frac{1}{F_1(x)}\right)^{1+s} \quad \text{and} \quad F_1(x) = F\left(\frac{1}{p}, \frac{1}{p}; 1 + \frac{1}{p}; x^p\right).$$

For (2), clearly f_2 is increasing for $m \geq 0$. For the case when $m \in [-1,0)$, we define

$$h_2(x) = \left(\frac{x}{\operatorname{artanh}_p x)}\right)^s \frac{1}{1-x^p}, \quad s \in (0,1].$$

Differentiating with respect to x, we get

$$h_2'(x) = \frac{(F_2(x))^{-(1+s)}\left((px^p - sx^p + s)\ F_2(x) - s\right)}{x\left(x^p - 1\right)^2} > 0,$$

where $F_2(x) = F\left(1, \frac{1}{p}; 1 + \frac{1}{p}; x^p\right)$.

For (3), the proof for the case when $m \geq 0$ follows similarly from Lemma 2.1. For the case $m \in [-1,0)$, let

$$h_3(x) = \left(\frac{\operatorname{arctan}_p x}{x}\right)^{-s} \frac{d}{dx}(\operatorname{arctan}_p x), \quad s \in (0,1].$$

We have

$$\begin{aligned} h_3'(x) &= \frac{F_3(x)^{-(1+s)}}{r(1+r^p)^2}((s+s\,r^p-p\,r^p)F_3(x)-s)\\ &< \frac{F_3(x)^{-(1+s)}}{r(1+r^p)^2}((s+s\,r^p-s\,r^p)F_3(x)-s)\\ &= -\frac{s\,F_3(x)^{-(1+s)}}{r(1+r^p)^2}(1-F_3(x))<0, \end{aligned}$$

where $F_3(x)=F\left(1,\frac{1}{p};1+\frac{1}{p};-x^p\right)$

For (4), when $m\geq 0$, the proof follows from Lemma 2.1. For $m\in[-1,0)$, let

$$h_4(x)=\left(\frac{x}{\operatorname{arsinh}_p x}\right)^s\frac{1}{(1+x^p)^{1/p}},\quad s\in(0,1]\,.$$

We have

$$\begin{aligned} h_4'(x) &= \gamma((1-x^p)^{1/p}(s(1-x^p)-x^p)F_4(x)-s(1-x^p))\\ &< \gamma\left(s(1+x^p)\left(1+\frac{1}{p}\log(1+x^p)\right)-s(1+x^p)-x^p\left(1+\frac{1}{1+p}\log(1+x^p)\right)\right)\\ &= \frac{\gamma}{p(1+p)}(s(1+x^p)(1+p)\log(1+x^p)-p(1+p)x^p-p\,x^p\log(1+x^p))\\ &< 0, \end{aligned}$$

by Lemma 2.2(4), where

$$\gamma=\frac{(1-x^p)^{-(1+2/p)}}{x}\left(\frac{1}{F_4(x)}\right)^{1+s}\quad\text{and}\quad F_4(x)=F\left(\frac{1}{p},\frac{1}{p};1+\frac{1}{p};-x^p\right).$$

□

2.5**. Proof of the Theorem 1.1.** Let $0<x<y<1$, and $u=((x^t+y^t)/2)^{1/t}>x$. We denote $\arcsin(x),\operatorname{artanh}(x),\arctan(x),\operatorname{arsinh}(x)$ by $g_i(x)$, $i=1,2\dots4$ respectively, and define

$$g(x)=g_i(u)^t-\frac{g_i(x)^t+g_i(y)^t}{2}.$$

Differentiating with respect to x, we get $du/dx=(1/2)(x/u)^{t-1}$ and

$$\begin{aligned} g'(x) &= \frac{1}{2}\,t\,g_i(x)^{t-1}\frac{d}{dx}(g_i(u))\left(\frac{x}{u}\right)^{t-1}-\frac{1}{2}\,t\,g_i(x)^{t-1}\frac{d}{dx}(g_i(x))\\ &= \frac{t}{2}x^{t-1}(f_i(u)-f_i(x)), \end{aligned}$$

where

$$f_i(x)=\left(\frac{g_i(x)}{x}\right)^{t-1}\frac{d}{dx}(g_i(x)),\ i=1,2\dots4.$$

By Lemma 2.4 g' is positive and negative for $f_{i=1,2}$ and $f_{i=3,4}$, respectively. This implies that

$$g(x) < (>) g(y) = 0,$$

for $g_{i=1,2}$ and $g_{i=3,4}$, respectively. The case when $t = 0$ follows from Lemma 2.3. This completes the proof.

2.6. **Lemma.** *For $p > 1$ and $s \in (0,1)$, the function*

$$f(p) = \left(\frac{\pi_p}{p}\right)^{-s} \frac{(p - \pi\cot(\pi/p))\csc(\pi/p)}{p^3}$$

is decreasing in $p \in (1,\infty)$,

Proof. We have

$$f'(p) = \xi\left[2p^2(1-s) + \pi^2(1-s)\cot^2\left(\frac{\pi}{p}\right) - \pi p(4-3s)\cot\left(\frac{\pi}{p}\right) + \pi^2\csc^2\left(\frac{\pi}{p}\right)\right],$$

which is negative, where

$$\xi = -\frac{(2\pi)^{-s}}{p^3}\left(\frac{\csc(\pi/2)}{p^2}\right)^{1-s}.$$

□

2.7. **Lemma.** [K, Thm 2, p.151] *Let $J \subset \mathbb{R}$ be an open interval, and let $f : J \to \mathbb{R}$ be strictly monotonic function. Let $f^{-1} : f(J) \to J$ be the inverse to f then*

(1) *if f is convex and increasing, then f^{-1} is concave,*
(2) *if f is convex and decreasing, then f^{-1} is convex,*
(3) *if f is concave and increasing, then f^{-1} is convex,*
(4) *if f is concave and decreasing, then f^{-1} is concave.*

2.8. **Lemma.** *For $m \geq 1$, $p > 1$ and $x \in (0,1)$, the following functions*

(1) $h_1(x) = \left(\frac{\sin_p x}{x}\right)^{m-1}\frac{d}{dx}(\sin_p x)$,
(2) $h_2(x) = \left(\frac{\tanh_p x}{x}\right)^{m-1}\frac{d}{dx}(\tanh_p x)$,
are decreasing in x, and
(3) $h_3(x) = \left(\frac{\cos_p x}{x}\right)^{m-1}\frac{d}{dx}(\cos_p x)$,
(4) $h_4(x) = \left(\frac{\tan_p x}{x}\right)^{m-1}\frac{d}{dx}(\tan_p x)$,
(5) $h_5(x) = \left(\frac{\sinh_p x}{x}\right)^{m-1}\frac{d}{dx}(\sinh_p x)$,

are increasing in x.

Proof. Let $f(x) = \mathrm{arcsin}_p\, x$, $x \in (0,1)$. We get

$$f'(x) = \frac{1}{(1-x^p)^{1/p}}\,,$$

which is positive and increasing, hence f is convex. Clearly $\sin_p\, x$ is increasing, and by Lemma 2.7 is concave, this implies that $\frac{d}{dx}\sin_p\, x$ is decreasing, and $(\sin_p x)/x$ is decreasing also by Lemma 2.1. Similarly we get that $\frac{d}{dx}\tanh_p\, x$ is decreasing and $\frac{d}{dx}\cos_p\, x$, $\frac{d}{dx}\tan_p\, x$, $\frac{d}{dx}\sinh_p\, x$ are increasing, and the rest of proof follows from Lemma 2.1. ☐

2.9**. Proof of the Theorem 1.2.** The proof is similar to the proof of Theorem 1.1 and follows from Lemma 2.8.

2.10. **Proposition.** *For $p,q>1$ and $t<1$, we have*

$$\pi_{M_t(p,q)} \le M_t(\pi_p,\pi_q)\,.$$

Proof. Let $1<p<q<\infty$, and $w = ((p^t+q^t)/2)^{1/t} > p$. We define

$$g(p) = (\pi_p)^t - \frac{(\pi_p)^t + (\pi_q)^t}{2}.$$

Differentiating with respect to p, we get $dw/dp = (1/2)(p/w)^{t-1}$ and

$$\begin{aligned} g'(p) &= \frac{1}{2}\,t\,(\pi_p)^{t-1}\frac{d}{dx}(\pi_w)\left(\frac{p}{w}\right)^{t-1} - \frac{1}{2}\,t\,(\pi_p)^{t-1}\frac{d}{dx}(\pi_p) \\ &= \frac{t}{2}p^{t-1}(f(w)-f(p)), \end{aligned}$$

where

$$f(p) = \left(\frac{\pi_p}{p}\right)^{t-1}\frac{d}{dp}\pi_p\,.$$

Clearly π_p is decreasing, hence $(\pi_p/p)^{t-1}$ is increasing for $t<1$ and $d/dp(\pi_p)$ is increasing by the proof of Lemma [BV1, Lemma 3.6]. This implies that $f(p)$ is increasing, and it follows that g is increasing. Hence $g(p)<g(q)=0$. The case when $t=0$ follows from Lemma 2.3(5). This completes the proof. ☐

The following lemma follows immediately from Lemma 2.8.

2.11. **Lemma.** *For $p>1$ and $r,s\in(0,1)$ with $r\le s$, we have*

(1) $\dfrac{\sin_p r}{r} \ge \dfrac{\sin_p s}{s}$,

(2) $\dfrac{\cos_p r}{r} \ge \dfrac{\cos_p s}{s}$,

(3) $\dfrac{\tan_p r}{r} \le \dfrac{\tan_p s}{s}$,

(4) $\dfrac{\sinh_p r}{r} \le \dfrac{\sinh_p s}{s}$,

$$\frac{\tanh_p r}{r} \geq \frac{\tanh_p s}{s}. \tag{5}$$

References

[AS] M. Abramowitz and I. Stegun, eds.: *Handbook of mathematical functions with formulas, graphs and mathematical tables.* National Bureau of Standards, 1964 (Russian translation, Nauka 1979).

[AVV1] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Conformal invariants, inequalities and quasiconformal maps.* J. Wiley, 1997, 505 pp.

[AVV2] G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen: *Generalized convexity and inequalities.* J. Math. Anal. Appl. 335 (2007), 1294–1308.

[BV1] B. A. Bhayo and M. Vuorinen: *Inequalities for eigenfunctions of the p-Laplacian.* January 2011, 23 pp. arXiv math.CA 1101.3911.

[BV2] B. A. Bhayo and M. Vuorinen: *On generalized trigonometric functions with two parameters.* J. Approx. Theory 164 (2012) 1415–1426. arXiv:1112.0483v.

[BEM1] R. J. Biezuner, G. Ercole, and E. M. Martins: *Computing the first eigenvalue of the p-Laplacian via the inverse power method.* J. Funct. Anal. 257 (2009), no. 1, 243–270.

[BEM2] R. J. Biezuner, G. Ercole, and E. M. Martins: *Computing the* $\sin_p$ *function via the inverse power method.* arXiv:1011.3486[math.CA].

[BE] P. J. Bushell and D. E. Edmunds: *Remarks on generalised trigonometric functions.* Rocky Mountain J. Math. Volume 42, Number 1 (2012), 25–57.

[EGL] D. E. Edmunds, P. Gurka, and J. Lang: *Properties of generalized trigonometric functions.* J. Approx. Theory 164 (2012) 47–56, doi:10.1016/j.jat.2011.09.004.

[DM] P. Drábek and R. Manásevich: *On the closed solution to some $p-$Laplacian nonhomogeneous eigenvalue problems.* Diff. and Int. Eqns. 12 (1999), 723–740.

[K] M. Kuczma: *An introduction to the theory of functional equations and inequalities.* Cauchy's equation and Jensen's inequality. With a Polish summary. Prace Naukowe Uniwersytetu Śląskiego w Katowicach [Scientific Publications of the University of Silesia], 489. Uniwersytet Śląski, Katowice; Państwowe Wydawnictwo Naukowe (PWN), Warsaw, 1985. 523 pp. ISBN: 83-01-05508-1.

[LE] J. Lang and D.E. Edmunds: *Eigenvalues, Embeddings and Generalised Trigonometric Functions.* Lecture Notes in Mathematics 2016, Springer-Verlag, 2011.

[L] P. Lindqvist: *Some remarkable sine and cosine functions.* Ricerche di Matematica, Vol. XLIV (1995), 269–290.

[LP] P. Lindqvist and J. Peetre: *p-arclength of the q-circle.* The Mathematics Student, Vol. 72, Nos. 1-4 (2003), 139–145.

[R] H. Ruskeepää: *Mathematica® Navigator.* 3rd ed. Academic Press, 2009.

[T] S. Takeuchi: *Generalized Jacobian elliptic functions and their application to bifurcation problems associated with p-Laplacian.* J. Math. Anal. 2011, doi:10.1016/j.jmaa.2011.06.063.

E-mail address: `barbha@utu.fi`

Department of Mathematics & Statistics, University of Turku, FI-20014 Turku, Finland
E-mail address: `vuorinen@utu.fi`